

Trees Nutrition Series

(Summary Sheet)



WSFNR10-32

December 2010

zn

element symbol

ZINC

by Dr. Kim D. Coder, Professor of Tree Biology & Health Care
Warnell School of Forestry & Natural Resources, University of Georgia

element number	30
element family type	METALS
normal form of pure element	SOLID METAL
at biological temperatures	
average rounded atomic weight	65
number of native isotopes	4
concentration group	DEKA-ELEMENT
element concentration in tree (ppm)	38
element proportion in tree	85
(carbon & oxygen levels = 450,000)	
element concentration rank in tree	13
(carbon & oxygen rank = 1)	
relative tree concentration	>
(compared to element in Earth's crust)	
different chemical oxidation states	1
most stable chemical oxidation state	2
oxidation states within a biologic compound	+2
oxidation states as a biologic active center	+2
total oxidation state range in biologics	1

among tree essential elements --	
relative atomic radius	LARGE
relative ionic radius	MEDIUM
relative first ionization energy	MEDIUM
relative atomic density	HIGH
other element family members (*toxic)	Cd*, Hg*
most commonly available tree form	Zn⁺²
(form in bold dominant)	
solubility of element's compounds --	
Zn⁺⁺ insoluble	= O⁻, S⁻, OH⁻, CO₃⁻
Zn⁺⁺ soluble	= NO₃⁻, SO₄⁻, C₂H₃O₂⁻

Coder Element Interaction Matrix for Trees (CEIMT) Values

(+ = positive or synergistic; - = negative or antagonistic)

B	Ca	Cl	Co	Cu	Fe	K	Mg	Mn	Mo	N _a	N _n	Ni	P	S	Si	Zn
+-	-	O	-	-	-	-	+-	-	O	-	-	-	-	-	O	X

In compliance with federal law, including the provisions of Title IX of the Education Amendments of 1972, Title VI of the Civil Rights Act of 1964, Sections 503 and 504 of the Rehabilitation Act of 1973, and the Americans with Disabilities Act of 1990, the University of Georgia does not discriminate on the basis of race, sex, religion, color, national or ethnic origin, age, disability, or military service in its administration of educational policies, programs, or activities; its admissions policies; scholarship and loan programs; athletic or other University-administered programs; or employment. In addition, the University does not discriminate on the basis of sexual orientation consistent with the University non-discrimination policy. Inquiries or complaints should be directed to the director of the Equal Opportunity Office, Peabody Hall, 290 South Jackson Street, University of Georgia, Athens, GA 30602. Telephone 706-542-7912 (V/TDD). Fax 706-542-2822. AN EQUAL OPPORTUNITY / AFFIRMATIVE ACTION INSTITUTION.

Zinc (Zn) is a hard, brittle, bluish-silver metal resistant to corrosion. Zinc can exist in ten isotopes, five stable, the rest all short-lived. It was known in the 1200s in India and later identified and named from German for “tin.” It is used for galvanizing steel and in batteries, coins, castings, paints, sunscreen, photocopiers, and cathode ray tubes.

Zinc is a required metal in trees. Zinc is divalent (+2) metal cation but unlike most of the other metals, does not in use undergo valence changes (i.e. no oxidation / reduction cycles). There are many zinc using or zinc activated enzymes in trees. Zinc function to activate proteins sometimes as the active site and sometime as a structural or conformational component. Many time zinc is seen cross-linking sulfur in proteins.

Zinc is required in trees for the proper transcription of DNA and gene expression. It is a key component in photosynthetic enzymes. Zinc is required for growth regulator (auxin) synthesis and for combining amino acids into proteins. Under anaerobic conditions, zinc helps detoxify alcohol accumulations. In soils, zinc at low to neutral pH is found in the form Zn^{2+} and at high pH is found in the form of $ZnOH^+$. High pH (>8.2) tends to generate insoluble zinc ($ZnCO_3$) and produce zinc deficiencies in trees. Figure 1. Figure 2.

Zinc deficiency in trees is first seen as leaves darkening and taking on a blue-green color which fades into a general yellowing. Leaves become stunted with a mottled appearance between the veins. Leaves eventually become distorted and die. Tree shoots become stunted with internodes not expanding. Shoots become distorted and die. Roots tend to exude gums and resins, and stop growth. Zinc deficiency is common in highly weathered and calcium rich soils with pH >8.2 where zinc becomes insoluble. In organic soils, or soils with a large amount of composted organic matter, zinc tends to become bound up and unavailable.

As zinc becomes more deficient, more phosphorus is taken up by trees. Zinc competes with nickel for transport and activation sites generating zinc deficiencies when nickel concentrations are too great. High concentrations of zinc suppresses potassium, calcium, and magnesium. Under anaerobic conditions, or through enrichment, cobalt minimizes this effect.

Zinc is easily added to enrich tree sites with many effective and low cost products. Traditionally, zinc nitrate ($Zn(NO_3)_2$) as a 1% foliar application has been used to small trees and shrubs. In some cases and under some conditions, this foliar spray cause leaf damage. Using $ZnSO_4$ as a 0.18% solution with hydrated lime has been cited as preventing zinc damage to leaves as a foliar spray. Zinc has not been found to be effective as a trunk injection or implant. Mycorrhizae in trees tend to mitigate and protect trees from zinc toxicity impacts.

Tree Symptom Summary

Zinc performs two dominant roles in trees: 1) Part of several enzymes constituents; and, 2) Activator / modifier of several enzymes. Deficiency symptoms can quickly occur physiologically downstream from these points.

When deficient, zinc has been cited as generating the following symptoms:

tree part	primary symptom	element deficiency responsible
roots	stunted / damaged	Zn -- also B, Cl, Cu, Mn, N, Ni, P, K, S, Si
	gum exuded (exanthema)	Zn -- also Cu
shoots	stunted / damaged / killed	Zn -- also B, Ca, Cl, Cu, Fe, Mn, Mo, N, Ni, P, K, S
	gum exuded (exanthema)	Zn -- also Cu
young leaf	wilting	Zn -- also B, Cl, Cu, K, Mo
leaves	color – blue-green / dark	Zn -- also Cl, K, P
	color -- dark viens	Zn -- also Cu, Mn, P
	color – general chlorosis	Zn -- also B, Cl, Cu, Fe, K, Mg, Mo, Mn, Ni, S
	intervienal chlorosis / death	Zn -- also Fe, Mg, Mo, Mn, Ni, S
	stunted / distorted blades	Zn -- also B, Cl, Cu, K, Mg, Mn, Mo, N, N
whole tree	growth regulator disruption / dysfunction	Zn -- also Co

Zinc is considered an intermediate among elements for mobility within a tree (immobile rank 8th).

symptom tissue location & age	element mobility inside tree	causal elemental deficiency
new tissues dominant	immobile	Zn -- also B, Ca, Co, Cu, Fe, Mn, Ni, S
all tissues equally	mobile	Zn -- also Cl, Cu, K, Ni, N, P, Si
intermediate	mobile/immobile	Zn -- also Mn, Mo, S

At pH 8.2 to 10.0, zinc is poorly available or unavailable to trees.

Proper identification of the cause for toxicity or deficiency symptoms must, at the least, involve tissue analysis for deficiencies and soil testing for toxicities.

relative
zinc
uptake




Figure 1: Estimated impact of soil pH on relative zinc (Zn) uptake in percent.

$\frac{\text{available Zn}}{\text{total Zn}}$
(percent)




Figure 2: Estimated soil pH impact on tree available zinc (Zn) as a percent of total soil zinc (Zn) concentration.